

# Parameterized Complexity: Exponential Speed-Up for Planar Graph Problems

Jochen Alber<sup>1</sup> Henning Fernau<sup>\*,2</sup> Rolf Niedermeier<sup>3</sup>

*Wilhelm-Schickard-Institut für Informatik, Universität Tübingen, Sand 13, D-72076 Tübingen, Fed. Rep. of Germany*

---

## Abstract

We discuss general techniques, centered around the “Layerwise Separation Property” (LSP) of a planar graph problem, that allow to develop algorithms with running time  $c^{\sqrt{k}}|G|$ , given an instance  $G$  of a problem on planar graphs with parameter  $k$ . Problems having LSP include PLANAR VERTEX COVER, PLANAR INDEPENDENT SET, and PLANAR DOMINATING SET. Extensions of our speed-up technique to basically all fixed-parameter tractable planar graph problems are also exhibited. Moreover, we relate, e.g., the domination number or the vertex cover number, with the treewidth of a plane graph.

*Key words:* Planar graph problems, fixed-parameter tractability, parameterized complexity, tree decomposition, graph separators

---

## 1 Introduction

**Fixed-parameter tractability.** While many problems of practical interest tend to be intractable from a standard complexity-theoretic point of view, in many cases such problems have natural “structural” parameters, and practically relevant instances are often associated with “small” values of these

---

\* Corresponding author

*Email addresses:* `alber@informatik.uni-tuebingen.de` (Jochen Alber), `fernau@informatik.uni-tuebingen.de`, `fernau@cs.newcastle.edu.au` (Henning Fernau), `niedermr@informatik.uni-tuebingen.de` (Rolf Niedermeier).

<sup>1</sup> Supported by the Deutsche Forschungsgemeinschaft, research project PEAL (Parameterized complexity and Exact ALgorithms), NI 369/1.

<sup>2</sup> Also affiliated with the University of Newcastle, Australia.

<sup>3</sup> Supported by the Deutsche Forschungsgemeinschaft, junior research group PIAF (fixed-parameter algorithms), NI 369/4.

parameters. The notion of fixed-parameter tractability [22] tries to capture this intuition (see [21,25] for some recent surveys). This is done by taking into account solving algorithms that are exponential with respect to the parameter, but otherwise have polynomial time complexity. That is, on input instance  $(I, k)$  one terms a (parameterized) problem *fixed-parameter tractable* if it allows for a solving algorithm running in time  $f(k)n^{O(1)}$ , where  $f$  is an arbitrary function only depending on  $k$  and  $n = |I|$ . The associated complexity class is called FPT. As fixed-parameter tractability explicitly allows for exponential time complexity concerning the parameter, the pressing challenge is to keep the related “combinatorial explosion” as small as possible.

In particular, the question naturally arises how “small” we can make the function  $f(k)$  [8]. One direction in current research on parameterized complexity is to investigate problems with fixed-parameter algorithms of running time  $c^k n^{O(1)}$  and to try to get the constant  $c$  as small as possible (e.g., see [4] in the case of PLANAR DOMINATING SET). Getting small constant bases in the exponential factor  $f(k)$  is also our concern, but, even more importantly, our primary focus is on investigations to get functions  $f$  (asymptotically) growing as slowly as possible. Doing so, we provide a general framework for a broad class of problems, namely planar graph problems. We indicate necessary conditions for planar graph problems that imply FPT-algorithms with running time  $c^{\sqrt{k}} n^{O(1)}$  for constant  $c$ . Moreover, we discuss an extension of our technique which applies to basically all fixed-parameter tractable graph problems.

In this paper, we provide a general framework for NP-hard planar graph problems that allows us to go from typically time  $c^k n^{O(1)}$  algorithms to time  $c^{\sqrt{k}} n^{O(1)}$  algorithms (subsequently briefly denoted by “ $c^{\sqrt{k}}$ -algorithms”), meaning an exponential speed-up.<sup>4</sup>

**Planar graph problems.** Planar graphs build a natural and practically important graph class. Many problems that are NP-complete for general graphs (such as VERTEX COVER and DOMINATING SET) remain so when restricted to planar graphs. Many NP-complete graph problems are hard to approximate in general graphs, but this is no longer true for restricted graph classes as planar graphs. Lipton and Tarjan observed that their separator theorems can be used to derive approximation schemes for problems on planar graphs [37,38]. Later, Baker [10] improved on the constants in the running times of these approximation schemes, coming with the use of separator theorems, by introducing the notion of outerplanarity.

Alternatively, finding an “efficient” exact solution in “reasonable exponen-

---

<sup>4</sup> Actually, whenever we can construct a so-called problem kernel of polynomial size in polynomial time (which is often the case for parameterized problems), then we can replace the term  $c^{\sqrt{k}} n^{O(1)}$  by  $c^{\sqrt{k}} k^{O(1)} + n^{O(1)}$ .

tial time” is an interesting and promising research challenge. In particular, many graph problems fixed-parameter intractable for general graph classes become fixed-parameter tractable for planar graphs (cf. [22]). This paper can be seen as an outerplanarity/treewidth based approach to the design of parameterized algorithms with sublinear exponents on planar graphs, this way supplementing the graph separator based approach detailed in [6]. Again, the outerplanarity/treewidth based approach turns out to be superior to the separator based approach in many respects, especially regarding the estimates on the running time of the algorithms.

**Methodology.** In recent work, algorithms were presented that constructively produce a solution for PLANAR DOMINATING SET and related problems in time  $c^{\sqrt{k}}n$  [1,2]. To obtain these results, it was proven that the treewidth of a planar graph with a dominating set of size  $k$  is bounded by  $O(\sqrt{k})$  and that a corresponding tree decomposition can be found in time  $O(\sqrt{kn})$ . Building on that problem-specific work with its rather tailor-made approach for dominating sets, here we take a much broader perspective. This comes to light by the following two main points. Firstly, by introducing the so-called “Layerwise Separation Property” we provide an abstract, problem-independent tool for a general design technique for  $c^{\sqrt{k}}$ -algorithms. Secondly, based on this key notion we give two main ways how to finally obtain these algorithms, one based on tree decompositions and the other one based on a bounded outerplanarity approach. Both these approaches do have their pros and cons, which will be discussed later on. (Please refer to Section 2 for a schematic picture of our methodology.) Note that, even though algorithms based on tree decompositions are widely considered to be impractical, because finding a tree decomposition, in general, is much too time-consuming, our approaches provide—to our knowledge—first results where such decompositions of small width can be computed relatively quickly, hence, yielding efficient algorithms. The advantage of both approaches from a practitioner’s point of view clearly is that, since the algorithms developed here can be stated in a very general framework, only small parts have to be changed to adapt them to the concrete problem. In this sense, our work differs strongly from research directions, where running times of algorithms are improved in a very problem-specific manner (e.g., by extremely sophisticated case-distinctions as in the case of VERTEX COVER for general graphs [17,42]). For example, once one can show that a problem has the so-called “Layerwise Separation Property,” one can run a general algorithm which quickly computes a tree decomposition of guaranteed small width (independent of the concrete problem). In summary, the heart of our approach can roughly be sketched as follows: If...

- (1) ...one can show that a graph problem carries some nice properties (e.g., the Layerwise Separation Property) and
- (2) ...one can determine some corresponding “problem-parameters” for these

properties (e.g., the width and the size-factor of the Layerwise Separation Property);

then one gets an algorithm of running time  $O(c^{\sqrt{k}}n^{O(1)})$ , where we give concrete formulas on how to evaluate the constant  $c$  as a function of these problem-parameters.

A library containing implementations of various algorithms sketched in this paper is currently under development. It uses the LEDA package [40] for efficient data types and (graph) algorithms and the results obtained so far are encouraging [3].

**The impact of the paper.** A preliminary version of this paper appeared in [7]. This paper, together with the conference version [1] of [2], inspired lots of research, partly hunting for better constants in the algorithms, partly extending the work towards larger graph classes. We shortly describe these efforts in the following.

- In [19], Demaine *et al.* show how to extend basically the results of this paper to the classes of  $K_5$ -minor-free graphs and of  $K_{3,3}$ -minor-free graphs, at the expense of worsening the constants in the running times. Note that the planar graphs are both  $K_5$ -minor-free and  $K_{3,3}$ -minor-free. Fomin and Thilikos explored and characterized the largest class admitting parameterized algorithms for DOMINATING SET: it turns out to be the class of graphs having bounded *local* treewidth [27].
- Fomin and Thilikos develop in [26,28] a branchwidth based counterpart to our approach [1,2,7]. They derive constants which are slightly better than ours, at the cost of a (in our opinion) more complicated proof, heavily relying on the sophisticated machinery developed by Robertson and Seymour. Kanj and Perkovič [31] and, with some additional geometric arguments, Fernau and Juedes (ongoing work) showed that the outerplanarity approach is also competitive with the branchwidth approach in the case of PLANAR DOMINATING SET. Probably, in practice it will be case-dependent which of the methods turns out to be superior.
- There has also been some progress on other planar graph problems (in terms of the involved constants) which are not in the focus of this paper: here, we mention Kloks, Lee and Liu's [35] contribution on FACE COVER, as well as Fernau and Juedes (ongoing work).

## 2 Outline and overview

The approach of this paper is based on the so-called layer-model of a planar graph as introduced in Subsection 3.2. Using this model, one proceeds in two




Fig. 1. Road map of our methodology for planar graph problems.

phases.

In a first phase, one separates the graph in a particular way (“layerwise”). The key property of a graph problem to allow such an approach will be the so-called “Layerwise Separation Property.” The details hereof are presented in Section 4. It will be shown that such a property holds for quite a large class of graph problems including those which admit a linear problem kernel. This property assures that the planar graph can be separated nicely.

In a second phase, the problem is solved on the layerwisely separated graph. We present two independent ways to achieve this in Section 5. Either, using the separators to set up a tree decomposition of width  $O(\sqrt{k})$  and solving the problem using this tree decomposition; or using a combination of a trivial approach on the separators and some algorithms working on graphs of bounded outerplanarity (see [10]) for the partitioned rest graphs. Figure 1 gives a general overview of our methodology presented in the following two sections. As noted before, in the second phase (Section 5) we will describe two independent ways to solve the underlying graph problem on the layerwisely separated graph. Both the tree decomposition as well as the bounded outerplanarity approach do have their pros and cons, which is why we present both of them. As to the tree decomposition approach, its advantage is its greater generality (up to the tree decomposition it is the same for all graph problems). In particular, it is definitely easier to implement in practice, also due to its universality and mathematical elegance.

By way of contrast, as to the bounded outerplanarity approach, in some cases we obtain better (theoretical worst-case) time complexity bounds for our algorithms in comparison with the tree decomposition approach. Moreover, the space consumption is significantly smaller, because the tree decomposition ap-

proach works in its dynamic programming part with possibly large tables. To achieve this, however, we need more complicated formalism and more constraints concerning the underlying graph problems.

### 3 Basic definitions and preliminaries

We consider undirected graphs  $G = (V, E)$ ,  $V$  denoting the vertex set and  $E$  denoting the edge set. For clarity, sometimes we refer to  $V$  by  $V(G)$ . Let  $G[D]$  denote the subgraph induced by a vertex set  $D \subseteq V$ . We only consider simple (no double edges) graphs without self-loops. We study *planar* graphs, i.e., graphs that can be drawn in the plane without edge crossings. Let  $(G, \phi)$  denote a *plane* graph, i.e., a planar graph  $G$  together with an embedding  $\phi$ . A *face* of a plane graph is any topologically connected region surrounded by edges of the plane graph. The one unbounded face of a plane graph is called the *exterior face*.

A *parameterized graph problem* is a language consisting of tuples  $(G, k)$ , where  $G$  is a graph and  $k$  is an integer. A *parameterized graph problem on planar graphs* is a parameterized graph problem, where the graph  $G$  of an instance  $(G, k)$  is assumed to be a planar graph.<sup>5</sup> Among others, we study the following “graph numbers”:

- A *vertex cover*  $C$  of a graph  $G$  is a set of vertices such that every edge of  $G$  has at least one endpoint in  $C$ ; the size of a vertex cover set with a minimum number of vertices is denoted by  $vc(G)$ .
- An *independent set* of a graph  $G$  is a set of pairwise nonadjacent vertices; the size of an independent set with a maximum number of vertices is denoted by  $is(G)$ .
- A *dominating set*  $D$  of a graph  $G$  is a set of vertices such that each of the rest of the vertices in  $G$  has at least one neighbor in  $D$ ;  $ds(G)$  denotes the size of a dominating set with a minimum number of vertices.

The corresponding problems are denoted by (PLANAR) VERTEX COVER (P)VC, (PLANAR) INDEPENDENT SET (P)IS, and (PLANAR) DOMINATING SET (P)DS.

---

<sup>5</sup> If the instances of a parameterized graph problem on planar graphs were of the form  $((G, \phi), k)$ , where  $\phi$  is an embedding of  $G$ , we would speak of a *parameterized graph problem on plane graphs*. In our setting, the planar graph  $G$  need not be given with an embedding.

### 3.1 Linear problem kernels

Reduction to problem kernel is a core technique for the development of fixed-parameter algorithms (see [22]). The idea behind is to cut off the “easy parts” of a given problem instance such that only the “hard kernel” of the problem remains where, then, e.g., exhaustive search can be applied (with reduced costs).

More formally, reduction to problem kernel here is defined as follows.

**Definition 1** *Let  $\mathcal{L}$  be a parameterized problem, i.e.,  $\mathcal{L}$  consists of pairs  $(I, k)$ , where problem instance  $I$  has a solution of size  $k$  (the parameter).<sup>6</sup> Reduction to problem kernel then means to replace the instance  $(I, k)$  by a “reduced” instance  $(I', k')$  (which we call the problem kernel) such that*

$$k' \leq d \cdot k, \quad |I'| \leq q(k')$$

*with constant  $d$ , polynomial  $q$ , and*

$$(I, k) \in \mathcal{L} \text{ iff } (I', k') \in \mathcal{L}.$$

*Furthermore, we require that the reduction from  $(I, k)$  to  $(I', k')$  (that we call kernelization) is computable in polynomial time  $T_K(|I|, k)$ , the subscript referring to kernelization time.*

Having constructed a size  $k^{O(1)}$  problem kernel in time  $n^{O(1)}$ , one can improve the time complexity  $f(k)n^{O(1)}$  of a fixed-parameter algorithm to  $f(k)k^{O(1)} + n^{O(1)}$ . Subsequently, our focus is on decreasing  $f(k)$ , and we do not always refer to this simple fact. Often, the best one can hope for the problem kernel is size linear in  $k$ , a so-called *linear problem kernel*.

**Example 2** *We discuss kernelizations for (P)VC, PIS, and PDS:*

**(P)VC** *Using a theorem of Nemhauser and Trotter [41], (also cf. [11,33,43]), Chen et al. [17] observed a problem kernel of size  $2k$  for (general) VC. According to the current state of knowledge, this is the best one could hope for because a problem kernel of size  $(2-\epsilon)k$  with constant  $\epsilon > 0$  would imply a factor  $2-\epsilon$  polynomial time approximation algorithm for VC, which would mean a major breakthrough in approximation algorithms for VC [29]. As regards PLANAR VERTEX COVER, however, a smaller problem kernel might be achievable.*

**PIS** *Due to the four color theorem for planar graphs and the corresponding algorithm generating a four coloring [44], it is easy to see that PIS has a*

<sup>6</sup> In this paper, we assume the parameter to be a positive integer, although, in general, it might also be from an arbitrary language (e.g., being a subgraph).

problem kernel of size  $4k$ . If we are looking for an independent set of size  $k \leq n/4$ , then the four coloring algorithm basically does the job. Otherwise, we know that  $k > n/4$ , i.e.,  $n < 4k$ , which gives a linear problem kernel.<sup>7</sup>

**PDS** The derivation of a problem kernel of size upperbounded by  $335k$  vertices for PDS is far from trivial [5].

Besides the positive effect of reducing the input size significantly by having small problem kernels, and all the already obvious consequences of that, this paper gives further justification, in particular, for the importance of linear problem kernels. The point is that once we have a linear problem kernel, it is fairly easy to get  $c^{\sqrt{k}}$ -algorithms for these problems based upon the famous planar separator theorem [37,38], see [6]. We will show alternative, more efficient ways (without using the planar separator theorem) of how to make use of linear problem kernels in a generic way in order to obtain  $c^{\sqrt{k}}$ -algorithms for planar graph problems, based on the ideas of Baker [10].

### 3.2 Tree decomposition and layer decomposition

In this subsection, we briefly introduce two sorts of decompositions for a graph  $G$ , which are a fundamental basis for our approaches.

**Definition 3** A tree decomposition of a graph  $G = (V, E)$  is a pair  $\langle \{X_i \mid i \in I\}, T \rangle$ , where  $X_i \subseteq V$  is called a bag and  $T$  is a tree with the elements of  $I$  as nodes, such that the following hold:

- (1)  $\bigcup_{i \in I} X_i = V$ ;
- (2) for every edge  $\{u, v\} \in E$ , there is an  $i \in I$  such that  $\{u, v\} \subseteq X_i$ ;
- (3) for all  $i, j, k \in I$ , if  $j$  lies on the path between  $i$  and  $k$  in  $T$ , then  $X_i \cap X_k \subseteq X_j$ .

The width of  $\langle \{X_i \mid i \in I\}, T \rangle$  is  $\max\{|X_i| \mid i \in I\} - 1$ . The treewidth  $\text{tw}(G)$  of  $G$  is the minimum  $\ell$  such that  $G$  has a tree decomposition of width  $\ell$ .

Details on tree decompositions can be found in [13,14,34].

The vertices of a planar graph  $G$  can be decomposed according to the level of the “layer” in which they appear in an embedding  $\phi$  [2,10].

**Definition 4** Let  $(G = (V, E), \phi)$  be a plane graph.

---

<sup>7</sup> Hence, it appears to be reasonable to look at the parameterized problem that asks for an independent set of size  $n/4 + k$  in the spirit of “parameterizing above guaranteed values” (cf. [39]).



- a) The layer decomposition of  $(G, \phi)$  is a disjoint partition of the vertex set  $V$  into sets  $L_1, \dots, L_r$ , which are recursively defined as follows:
- $L_1$  is the set of vertices on the exterior face of  $G$ .
  - $L_i$  is the set of vertices on the exterior face of  $G[V - \bigcup_{j=1}^{i-1} L_j]$  for all  $i = 2, \dots, r$ .

We will denote the layer decomposition of  $(G, \phi)$  by

$$\mathcal{L}(G, \phi) := (L_1, \dots, L_r).$$

- b) The set  $L_i$  is called the  $i$ th layer of  $(G, \phi)$ .  
 c) The (uniquely defined) number  $r$  of different layers is called the outerplanarity of  $(G, \phi)$ , denoted by  $\text{out}(G, \phi) := r$ .  
 d) We define  $\text{out}(G)$  to be the smallest outerplanarity possible among all plane embeddings, i.e., minimizing over all plane embeddings  $\phi$  of  $G$  we set

$$\text{out}(G) := \min_{\phi} \text{out}(G, \phi).$$

Due to technical reasons, for a layer-decomposition  $\mathcal{L}(G, \phi) := (L_1, \dots, L_r)$ , we set  $L_i := \emptyset$  for all indices  $i < 1$  and  $i > r$ .

Computing the layers of a plane graph can be done efficiently:<sup>8</sup>

**Proposition 5** *Let  $(G = (V, E), \phi)$  be a plane graph. The layer decomposition  $\mathcal{L}(G, \phi) = (L_1, \dots, L_r)$  can be computed in time  $O(|V|)$ .*

### 3.3 Algorithms based on separators in graphs

One of the most useful algorithmic techniques for solving computational problems is divide-and-conquer. To apply this method to planar graphs, we need graph separators and related notions.

To simplify notation, we write  $A + B$  for the disjoint union of two sets  $A$  and  $B$ . Graph separators are defined as follows.

**Definition 6** *Let  $G = (V, E)$  be an undirected graph. A separator  $S \subseteq V$  of  $G$  partitions  $V$  into two sets  $A_1$  and  $A_2$  such that*

- $V = A_1 + S + A_2$  and
- no edge joins vertices in  $A_1$  and  $A_2$ .

The triple  $(A_1, S, A_2)$  is called a separation of  $G$ , and the graphs  $G[A_i]$  are called the graph chunks of the separation. Given a separation  $(A_1, S, A_2)$ , we use the shorthands  $\delta A_i := A_i + S$  for  $i = 1, 2$ .

<sup>8</sup> For a detailed description of the algorithm, see [2, Section 2.3].

In general,  $A_1$ ,  $A_2$ , and  $S$  will be non-empty. In order to cover boundary cases in some considerations below, we did not put this into the separator definition. We will also consider  $X$ - $Y$ -separators with  $X, Y \subseteq V$ : such a separator cuts every path from  $X$  to  $Y$  in  $G$ .

Already Lipton and Tarjan [37,38] used their famous separator theorem in order to design algorithms with running time of  $O(c^{\sqrt{n}})$  for certain planar graph problems, e.g., for glueable graph problems (for a formal definition see [6]). This naturally implies that, in the case of parameterized planar graph problems for which a linear kernel is known, algorithms with running time  $O(c^{\sqrt{k}} + T_K(n, k))$  can be derived, where  $T_K(n, k)$  is the time to construct the problem kernel of an  $n$ -vertex input graph. As worked out in [6], it is possible to get an

$$O(2^{1.97\sqrt{dk}/(1-\sqrt{2/3+\epsilon})} + T_K(n, k))$$

algorithm for glueable planar graph problems with problem kernel of size  $dk$ , where  $\epsilon \in (0, 1/3)$  can be chosen freely.<sup>9</sup> Even if we can assume  $\epsilon = 0$  (this is feasible due to the nice properties of VERTEX COVER) we obtain an exponential growth of  $c^{\sqrt{k}}$  with  $c \approx 2^{15.19} \approx 37381$  for PLANAR VERTEX COVER, where  $d = 2$  is known from [17,41]. As detailed in [6], we can obtain divide-and-conquer algorithms with an exponential growth of  $2^{13.07\sqrt{k}} \approx 8564^{\sqrt{k}}$  by using separator theorems in a more clever way.

We will see algorithms with much better constants in this paper. For example, in the case of PVC, we obtain an algorithm with an exponential growth of  $2^{2\sqrt{6k}} \approx 2^{4.90\sqrt{k}} \approx 29.84^{\sqrt{k}}$ . This is only slightly worse than the recently obtained upperbound of  $2^{4.5\sqrt{k}}$  due to Fomin and Thilikos [26] based on a branchwidth approach. Compared to the separator based approach, the approach pursued in this paper relies on weaker assumptions. Thus, in some cases we may drop requirements such as linear problem kernels by replacing it with the so-called “Layerwise Separation Property,” a seemingly less restrictive demand.

#### 4 Phase 1: Layerwise separation

In this section, we exploit the layer-structure of a plane graph in order to gain a “nice” separation of the graph. It is important that a “yes”-instance  $(G, k)$  (where  $G$  is a plane graph) of the graph problem  $\mathcal{G}$  admits a so-called “layerwise separation” of small size. By this, we mean, roughly speaking, a separation of the plane graph  $G$  (i.e., a collection of separators for  $G$ ), such

---

<sup>9</sup> The constants 1.97 and 2/3 appearing in the exponent of the running time result from the best known  $\sqrt{\cdot}$ -separator theorem due to Djidjev and Venkatesan [20].

that each separator is contained in the union of constantly many subsequent layers (see conditions 1 and 2 of the following definition). For (fixed-parameter) algorithmic purposes, it will be important that the corresponding separators are “small” (see condition 3 of the definition).

**Definition 7** Let  $(G = (V, E), \phi)$  be a plane graph of outerplanarity  $r := \text{out}(G, \phi)$ , and let  $\mathcal{L}(G, \phi) = (L_1, \dots, L_r)$  be its layer decomposition. A layerwise separation of width  $w$  and size  $s$  of  $(G, \phi)$  is a sequence  $(S_1, \dots, S_r)$  of subsets of  $V$ , with the properties that, for  $i = 1, \dots, r$ :<sup>10</sup>

- (1)  $S_i \subseteq \bigcup_{j=i}^{i+(w-1)} L_j$ ,
- (2)  $S_i$  is an  $L_{i-1}$ - $L_{i+w}$  separator, and
- (3)  $\sum_{j=1}^r |S_j| \leq s$ .

The crucial property that makes the algorithms developed in this paper work is what we call the “Layerwise Separation Property.”

**Definition 8** A parameterized problem  $\mathcal{G}$  for planar graphs is said to have the Layerwise Separation Property (abbreviated by: LSP) of width  $w$  and size-factor  $d$  if for each  $(G, k) \in \mathcal{G}$  and every planar embedding  $\phi$  of  $G$ , the plane graph  $(G, \phi)$  admits a layerwise separation of width  $w$  and size  $dk$ .

#### 4.1 How can layerwise separations be obtained?

The LSP can be shown directly for many parameterized graph problems.

- Example 9** (1) Consider PVC. Here, we get constants  $w = 2$  and  $d = 2$ . In fact, for  $(G, k) \in \text{VERTEX COVER}$  (and any planar embedding  $\phi$  of  $G$ ) with a “witnessing” vertex cover  $V'$  of size  $k$ , the sets  $S_i := (L_i \cup L_{i+1}) \cap V'$  form a layerwise separation, given the layer decomposition  $\mathcal{L}(G, \phi) = (L_1, \dots, L_r)$ .
- (2) In [2], the non-trivial fact is proven that for PDS this property holds, yielding constants  $w = 3$  and  $d = 51$ .<sup>11</sup>

Fenau and Juedes (ongoing work) have shown that all problems describable by formulas from PLANAR TMIN<sub>1</sub> (as defined in [32]) satisfy LSP. This class includes PLANAR RED-BLUE DOMINATING SET, FACE COVER, and PLANAR EDGE DOMINATING SET.

<sup>10</sup> By default, we let  $S_i := \emptyset$  for all  $i < 1$  and  $i > r$ .

<sup>11</sup> Note that in the case of PLANAR DOMINATING SET a construction of this form (i.e., obtaining the separators  $S_i$  by intersecting a “witnessing” dominating set  $V'$  of  $G$  with a sequence of subsequent layers, e.g,  $S_i := (L_{i-1} \cup L_i \cup L_{i+1}) \cap V'$ ), does not fulfill the conditions of a layerwise separation, since, in general,  $S_i$  need not be a separator.

A large class of parameterized graph problems for which the LSP holds is given whenever there exists a reduction to a linear problem kernel.

**Lemma 10** *Let  $\mathcal{G}$  be a parameterized problem for planar graphs that admits a problem kernel of size  $dk$ . Then, the parameterized problem  $\mathcal{G}'$  where each instance is replaced by its problem kernel has the LSP of width 1 and size-factor  $d$ .*

**PROOF.** Let  $(G', k') \in \mathcal{G}'$  with  $k' \leq dk$  be the problem kernel of  $(G, k) \in \mathcal{G}$ , and let  $\mathcal{L}(G', \phi') = (L'_1, \dots, L'_{r'})$  be the layer decomposition of  $(G', \phi')$  (where  $\phi'$  is any embedding). Let  $r' = \text{out}(G', \phi')$ . Observe that  $r' \leq \frac{dk}{3}$  since each layer has to consist of at least 3 vertices. Then, the sequence  $S_i := L'_i$  for  $i = 1, \dots, r'$  is a layerwise separation of width 1 and size  $dk$  of  $(G', \phi')$ .

Example 2 therefore immediately yields:

**Example 11** (1) *PVC has the LSP of width 1 and size-factor 2 (which is even better than what was shown in Example 9).*

(2) *PIS has the LSP of width 1 and size-factor 4 on the set of reduced instances.*

(3) *PIS has the LSP of width 1 and size-factor 335. Observe that these bounds are worse than the one mentioned in Example 9, since it is the size-factor that mainly influences the running time of our algorithms.*

#### 4.2 What are layerwise separations good for?

From a layerwise separation of small size (say bounded by  $O(k)$ ), we are able to choose a set of separators such that their size is bounded by  $O(\sqrt{k})$  and—at the same time—the subgraphs into which these separators cut the original graph have outerplanarity bounded by  $O(\sqrt{k})$ . In order to formalize such a choice of separators from a layerwise separation, we give the following definition.

**Definition 12** *Let  $(G = (V, E), \phi)$  be a plane graph with layer decomposition  $\mathcal{L}(G, \phi) = (L_1, \dots, L_r)$ . A partial layerwise separation of width  $w$  is a sequence  $\mathcal{S} = (S_1, \dots, S_q)$  such that there exist  $i_0 = 1 < i_1 < \dots < i_q < r = i_{q+1}$  such that for  $i = 1, \dots, q$ :*<sup>12</sup>

- (1)  $S_j \subseteq \bigcup_{\ell=i_j}^{i_{j+1}+(w-1)} L_\ell$ ,
- (2)  $i_j + w \leq i_{j+1}$  (so, the sets in  $\mathcal{S}$  are pairwise disjoint) and

<sup>12</sup> Again, by default, we set  $S_i := \emptyset$  for  $i < 1$  and  $i > q$ .

(3)  $S_j$  is a  $L_{i_{j-1}}-L_{i_j+w}$  separator.

The sequence  $\mathcal{C}_S = (G_0, \dots, G_q)$  with

$$G_j := G\left[\bigcup_{\ell=i_j}^{i_{j+1}+(w-1)} L_\ell - (S_j \cup S_{j+1})\right], \quad j = 0, \dots, q,$$

is called the sequence of graph chunks obtained by  $\mathcal{S}$ .

With this definition at hand, we can state the key result needed to establish the algorithms that will be presented in Section 5. The proof techniques applied show some similarity to Baker [10].

**Proposition 13** *Let  $(G = (V, E), \phi)$  be a plane graph that admits a layerwise separation of width  $w$  and size  $dk$ . Then, for every  $\psi \in \mathbb{R}_+$ , there exists a partial layerwise separation  $\mathcal{S}(\psi)$  of width  $w$  such that*

- (1)  $\sum_{S \in \mathcal{S}(\psi)} |S| \leq \psi \sqrt{dk}$  and<sup>13</sup>
- (2)  $\text{out}(H) \leq \frac{\sqrt{dk}}{\psi} + w$  for each graph chunk  $H$  in  $\mathcal{C}_{\mathcal{S}(\psi)}$ .

Moreover, there is an algorithm with running time  $O(\sqrt{kn})$  which, for a given  $\psi$ ,

- recognizes whether  $(G, \phi)$  admits a layerwise separation of width  $w$  and size  $dk$  and, if so, computes  $\mathcal{S}(\psi)$ ;
- computes a partial layerwise separation of width  $w$  that fulfills the conditions above.

**PROOF.** For  $m = 1, \dots, w$ , consider the integer sequences  $I_m = (m + jw)_{j=0}^{\lfloor r/w \rfloor - 1}$  and the corresponding sequences of separators  $\mathcal{S}_m = (S_i)_{i \in I_m}$ . Note that each  $\mathcal{S}_m$  is a sequence of pairwise disjoint separators. Since  $(S_1, \dots, S_r)$  is a layerwise separation of size  $dk$ , this implies that there exists a  $1 \leq m' \leq w$  with

$$\sum_{i \in I_{m'}} |S_i| \leq \frac{dk}{w}. \quad (1)$$

For a given  $\psi$ , let  $s := \psi \sqrt{dk}$ . Define  $\mathcal{S}(\psi)$  to be the subsequence of  $\mathcal{S}_{m'}$  such that  $|S| \leq s$  for all  $S \in \mathcal{S}(\psi)$ , and  $|S| > s$  for all  $S \in \mathcal{S}_{m'} - \mathcal{S}(\psi)$ . This yields condition 1. As to condition 2, suppose that  $\mathcal{S}(\psi) = (S_{i_1}, \dots, S_{i_q})$ . How many layers are two separators  $S_{i_j}$  and  $S_{i_{j+1}}$  apart? Herefore, note that the number of separators in  $\mathcal{S}_{m'}$  that appear between  $S_{i_j}$  and  $S_{i_{j+1}}$  is  $(i_{j+1} - i_j)/w$ . Since all of these separators have size greater than  $s$ , their number has to be bounded by  $dk/ws$ , see Equation (1). Therefore, we get  $i_{j+1} - i_j \leq \sqrt{dk}/\psi$  for

<sup>13</sup> Taking  $\sum$  instead of  $\max$  here is a proposal of Kanj and Perkovič, compare [7,31].

all  $j = 1, \dots, q - 1$ . Hence, the chunks  $G[(\bigcup_{\ell=i_j}^{i_{j+1}+w-1} L_\ell) - (S_{i_j} \cup S_{i_{j+1}})]$  have outerplanarity at most  $\sqrt{dk}/\psi + w$ .

The algorithm that computes a partial layerwise separation  $\tilde{\mathcal{S}}$  proceeds as follows: For given  $\psi$ , compute  $s := \psi\sqrt{dk}$ . Then, for  $j = 1, \dots, r - w$ , one checks whether the graph  $\tilde{G}_j(v_s, v_t)$  admits a  $v_s$ - $v_t$ -separator  $\tilde{S}_j$  of size at most  $s$ . Here,  $\tilde{G}_j(v_s, v_t)$  is the graph  $G[\bigcup_{\ell=j}^{j+(w-1)} L_\ell]$  with two further vertices  $v_s$  and  $v_t$  and edges from  $v_s$  to all vertices in  $L_j$  and from  $v_t$  to all vertices in  $L_{j+w-1}$ . The separator  $\tilde{S}_j$  can be computed in time  $O(s \cdot n)$  using techniques based on maximum flow (see [30] for details).

Let  $\tilde{\mathcal{S}} = (\tilde{S}_1, \dots, \tilde{S}_q)$  be the sequence of all separators of size at most  $s$  found in this manner.<sup>14</sup> Suppose that  $\tilde{S}_j \subseteq \bigcup_{\ell=i_j}^{i_{j+1}+(w-1)} L_\ell$  for some indices  $1 \leq i_1 < \dots < i_q \leq r$ . Note that, by the arguments given above, no two such separators can be more than  $\sqrt{dk}/\psi$  layers apart. Hence, if there was a  $j_0$  such that  $i_{j_0+1} - i_{j_0} > \sqrt{dk}/\psi$ , the algorithm exits and returns “no.” Otherwise,  $\tilde{\mathcal{S}}$  is a partial layerwise separation of width  $w$ .

In what follows, the positive real number  $\psi$  of Proposition 13 is also called *trade-off parameter*. This is because it allows us to optimize the trade-off between outerplanarity and separator size.

Proposition 13 will help construct a tree decomposition of treewidth  $\text{tw}(G) = O(\sqrt{k})$ , assuming that a layerwise separation of some constant width and size  $dk$  exists. Hence, for graph problems fulfilling this assumption and, moreover, allowing a  $\sigma^{\text{tw}(G)}n$  time algorithm for constant  $\sigma$  when the graph is given together with a tree decomposition, we obtain a solving algorithm with running time  $c^{\sqrt{k}}n$ . This aspect will be outlined in Subsection 5.1.

## 5 Phase 2: Algorithms on layerwisely separated graphs

After Phase 1, we are left with a set of disjoint (layerwise) separators of size  $O(\sqrt{k})$  separating the graph in components, each of which having outerplanarity bounded by  $O(\sqrt{k})$ . We now present two different ways how to obtain, in a second phase, a  $c^{\sqrt{k}}$ -algorithm that makes use of this separation. In both cases, there is the trade-off parameter  $\psi$  from Proposition 13 that can be used to optimize the running time of the resulting algorithms.

<sup>14</sup>Possibly, the separators  $\tilde{S}_j$  in  $\tilde{\mathcal{S}}$  found by the algorithm may differ from the separators in  $\mathcal{S}(\psi)$ .

### 5.1 Using tree decompositions

We use the concept of tree decompositions as, e.g., described in [13,34] (also cf. Subsection 3.2). We show how the existence of a layerwise separation of small size helps to constructively obtain a tree decomposition of small width. The following result can be found in [36, Table 2, page 550] or [14, Theorem 83]. The corresponding algorithm is outlined in [2].

**Proposition 14** *For a plane graph  $(G, \phi)$ , we have  $\text{tw}(G) \leq 3 \cdot \text{out}(G) - 1$ . Such a tree decomposition can be found in  $O(\text{out}(G) \cdot n)$  time.*

With this proposition at hand, we can prove our central result in this context.

**Theorem 15** *Let  $(G, \phi)$  be a plane graph that admits a layerwise separation of width  $w$  and size  $dk$ . Then, we have  $\text{tw}(G) \leq 2\sqrt{3dk} + (3w - 1)$ . Such a tree decomposition can be computed in time  $O(k^{3/2}n)$ .*

**PROOF.** By Proposition 13, for each  $\psi \in \mathbb{R}_+$ , there exists a partial layerwise separation  $\mathcal{S}(\psi) = (S_1, \dots, S_q)$  of width  $w$  with corresponding graph chunks  $\mathcal{C}_{\mathcal{S}(\psi)} = (G_0, \dots, G_q)$ , such that  $\sum_{S \in \mathcal{S}(\psi)} |S| \leq \psi\sqrt{dk}$  and  $\text{out}(G_i) \leq \frac{\sqrt{dk}}{\psi} + w$  for all  $i = 0, \dots, q$ . The algorithm that constructs a tree decomposition  $\mathfrak{X}_\psi$  is given as follows:

- Construct a tree decomposition  $\mathcal{X}_i$  of width at most  $3 \text{out}(G_i) - 1$  for each of the graphs  $G_i$  (using the algorithm from Proposition 14).
- Add  $S_i$  and  $S_{i+1}$  to every bag in  $\mathcal{X}_i$  ( $i = 0, \dots, q$ ).
- Let  $T_i$  be the tree of  $\mathcal{X}_i$ . Then, successively add an arbitrary connection between the trees  $T_i$  and  $T_{i+1}$  in order to obtain a tree  $T$ .

It is easy to show (see [1, Proposition 4]) that the tree  $T$ , together with the constructed bags, gives a tree decomposition of  $G$ . Clearly, its width  $\text{tw}(\mathfrak{X}_\psi)$  is upperbounded by

$$\begin{aligned} \text{tw}(\mathfrak{X}_\psi) &\leq \sum_{S \in \mathcal{S}(\psi)} |S| + \max_{i=0, \dots, q} \text{tw}(G_i) \\ &\leq \sum_{S \in \mathcal{S}(\psi)} |S| + 3(\max_{i=0, \dots, q} \text{out}(G_i)) - 1 \\ &\leq (\psi + 3/\psi)\sqrt{dk} + (3w - 1). \end{aligned}$$

This upper bound is minimized for  $\psi = \sqrt{3}$ .

By [9, Proposition 4.5], a graph  $G$  that has no  $K_h$  minor has treewidth bounded by  $h^{3/2}\sqrt{n}$ . In particular, this implies that a planar graph has treewidth bounded by  $11.2\sqrt{n}$ . In the case of the existence of linear problem kernels for a given graph problem  $\mathcal{G}$ , this method might be used in order to obtain  $c^{\sqrt{k}}$ -algorithms. From our results, we can derive upper bounds of the treewidth of a planar graph in terms of several graph specific numbers. As the reader may verify, these problem-specific treewidth bounds tend to outperform the numbers obtainable via [9, Proposition 4.5]. For example, Theorem 15 and Example 9 imply the following inequalities for a planar graph  $G$ , relating the treewidth with the vertex cover and dominating set number:

$$\begin{aligned} \text{tw}(G) &\leq 2\sqrt{6\text{vc}(G)} + 5, \quad \text{and} \\ \text{tw}(G) &\leq 6\sqrt{17\text{ds}(G)} + 8. \end{aligned}$$

Note that for general graphs, no relation of the form

$$\text{tw}(G) \leq f(\text{ds}(G)) \tag{2}$$

(for any function  $f$ ) holds; consider, e.g., the clique  $K_n$  with  $n$  vertices, where  $\text{tw}(K_n) = n - 1$ , but  $\text{ds}(K_n) = 1$ . Fomin and Thilikos have recently shown [27] that Eq. (2) holds for a graph  $G$  iff  $G$  has bounded local treewidth. For VC, only the linear relation

$$\text{tw}(G) \leq \text{vc}(G)$$

can be easily shown: Note that the complement of a vertex cover set  $C$  forms an independent set  $I$  in  $G$ . Hence, we can easily construct even a path decomposition by choosing  $|I|$  bags and making each bag consist of all vertices of  $C$  and exactly one element from  $I$  each time. This estimate is sharp (which becomes clear by, again, considering the graph  $K_n$ , where  $\text{vc}(K_n) = n - 1$ ).

Theorem 15 yields a  $c^{\sqrt{k}}$ -algorithm for certain graph problems:

**Theorem 16** *Let  $\mathcal{G}$  be a parameterized problem for planar graphs. Suppose that*

- (1)  $\mathcal{G}$  has the LSP of width  $w$  and size-factor  $d$ , and
- (2) there exists a time  $\sigma^\ell$  algorithm that decides  $(G, k) \in \mathcal{G}$ , if  $G$  is given together with a tree decomposition of width  $\ell$ .

*Then, there is an algorithm to decide  $(G, k) \in \mathcal{G}$  in time  $O(\sigma^{3w-1} \cdot 2^{\theta_1(\sigma, d)\sqrt{k}n})$ , where  $\theta_1(\sigma, d) = 2(\log \sigma)\sqrt{3d}$ .*



**PROOF.** Given an instance  $(G, k)$ , in linear time we can compute some planar embedding  $\phi$  of  $G$  (for details see [18]). In time  $O(\sqrt{kn})$  (see Proposition 13), we can check whether the plane graph  $(G, \phi)$  admits a layerwise separation of width  $w$  and size  $dk$ .

If so, the algorithm of Theorem 15 computes a tree decomposition of width at most  $2\sqrt{3dk} + (3w - 1)$ , and we can decide  $(G, k) \in \mathcal{G}$  by using the given tree decomposition algorithm in time  $O(\sigma^{2\sqrt{3dk}+(3w-1)}n)$ .

If  $(G, \phi)$  does not admit such a layerwise separation, we know that  $(G, k) \notin \mathcal{G}$ , by definition of the LSP.

**Example 17** *Going back to our running examples, it is well-known that VERTEX COVER and INDEPENDENT SET admit such a tree decomposition based algorithm with  $\sigma = 2$  and, in the case of DOMINATING SET, with  $\sigma = 4$ , as detailed in [2].*

- (1) *For PVS, by Example 11.1 Theorem 16 guarantees an  $O(2^{2\sqrt{6k}}n)$  algorithm for this problem.*
- (2) *For PIS, Example 11.2 and Theorem 16 yield an  $O(2^{4\sqrt{3k}}n)$  algorithm.*
- (3) *By Example 9.2, Theorem 16 improves on the result from [2] (where we got an  $O(4^{6\sqrt{34k}}n) \approx O(2^{69.98\sqrt{k}}n)$  algorithm), namely, getting an  $O(4^{6\sqrt{17k}}n) \approx O(2^{49.48\sqrt{k}}n)$  algorithm for PLANAR DOMINATING SET.*

In this subsection, we have seen that, for plane graphs, the notion of the LSP gives us a sufficient condition to upperbound the treewidth of the “yes”-instance graphs of a problem. Moreover, this property led to *fast* computations of the corresponding tree decompositions. All in all, we came up with algorithms of running time  $O(c^{\sqrt{k}}n)$  for a wide class of problems.

The next subsection aims to show similar results in a different context.

## 5.2 Using bounded outerplanarity

We now turn our attention to certain parameterized graph problems for which we know that a solving algorithm of linear running time on the class of graphs of bounded outerplanarity exists. This issue was addressed in [10]; several examples can be found therein. In this subsection, we introduce the so-called “select&verify” problems and examine how, in this context, the notion of the LSP will lead to  $c^{\sqrt{k}}$ -algorithms. Since this will be a purely separator based approach, we will restrict ourselves to parameterized graph problems that can be solved easily on separated graphs. We will introduce the notion of weakly glueable select&verify problems in a first paragraph and present the design of  $c^{\sqrt{k}}$ -algorithms for these problems afterwards (see Paragraph 5.2.2).

5.2.1 *Select&verify problems and weak glueability*

For the approach outlined in this section, we want to describe necessary properties for graph problems that allow for separator based algorithms. The notion of select&verify graph problems as introduced in the next paragraph is exactly the same as in [6]. There, one also finds the more restrictive concept of “glueability,” which is a property tailored for divide-and-conquer algorithms using graph separators. In our setting, however, we are interested in an iterative dynamic programming approach for which a weaker form of “glueability” is sufficient.

**Select&verify graph problems**

**Definition 18** *A set  $\mathcal{G}$  of tuples  $(G, k)$ ,  $G$  an undirected graph with vertex set  $V = \{v_1, \dots, v_n\}$  and  $k$  a positive real number, is called a select&verify (graph) problem if there exists a pair  $(P, \text{opt})$  with  $\text{opt} \in \{\min, \max\}$ , such that  $P$  is a function that assigns to  $G$  a polynomial time computable function of the form  $P_G = P_G^{\text{sel}} + P_G^{\text{ver}}$ , where  $P_G^{\text{sel}} : \{0, 1\}^n \rightarrow \mathbb{R}_+$ ,  $P_G^{\text{ver}} : \{0, 1\}^n \rightarrow \{0, \pm\infty\}$ , and*

$$(G, k) \in \mathcal{G} \quad \Leftrightarrow \quad \begin{cases} \text{opt}_{\vec{x} \in \{0,1\}^n} P_G(\vec{x}) \leq k & \text{if } \text{opt} = \min \\ \text{opt}_{\vec{x} \in \{0,1\}^n} P_G(\vec{x}) \geq k & \text{if } \text{opt} = \max. \end{cases}$$

For  $\vec{x} = (x_1, \dots, x_n) \in \{0, 1\}^n$  with  $P_G(\vec{x}) \leq k$  if  $\text{opt} = \min$  and with  $P_G(\vec{x}) \geq k$  if  $\text{opt} = \max$ , the vertex set selected by  $\vec{x}$  and verified by  $P_G$  is  $\{v_i \in V \mid x_i = 1, 1 \leq i \leq n\}$ . A vector  $\vec{x}$  is called admissible if  $P_G^{\text{ver}}(\vec{x}) = 0$ .

The intuition behind the term  $P = P^{\text{sel}} + P^{\text{ver}}$  is that the “selecting function”  $P^{\text{sel}}$  counts the size of the selected set of vertices and the “verifying function”  $P^{\text{ver}}$  verifies whether this choice of vertices is an admissible solution.

Observe that every select&verify graph problem that additionally admits a problem kernel of size  $p(k)$  is solvable in time  $O(2^{p(k)}p(k) + T_K(n, k))$ .

**Example 19** *We now give some examples for select&verify problems by specifying the function  $P_G = P_G^{\text{sel}} + P_G^{\text{ver}}$ . In all cases below, the “selecting function”  $P_G$  for a graph  $G = (V, E)$  will be*

$$P_G^{\text{sel}} = \sum_{v_i \in V} x_i.$$

Also, we use the convention that  $0 \cdot (\pm\infty) = 0$ .

(1) *In the case of VERTEX COVER, we have  $\text{opt} = \min$  and choose*

$$P_G^{\text{ver}}(\vec{x}) = \sum_{\{v_i, v_j\} \in E} \infty \cdot (1 - x_i)(1 - x_j),$$

where this sum brings  $P_G(\vec{x})$  to infinity whenever there is an uncovered edge. In addition,  $P_G(\vec{x}) \leq k$  then guarantees a vertex cover set of size at most  $k$ . Clearly,  $P_G$  is polynomial time computable.

- (2) Similarly, in the case of INDEPENDENT SET, let  $\text{opt} = \max$  and choose

$$P_G^{\text{ver}}(\vec{x}) = \sum_{\{v_i, v_j\} \in E} (-\infty) \cdot x_i \cdot x_j.$$

- (3) DOMINATING SET is another example for a select&verify graph problem. Here, for  $G = (V, E)$ , we have

$$P_G^{\text{ver}}(\vec{x}) = \sum_{v_i \in V} (\infty \cdot (1 - x_i) \cdot \prod_{\{v_i, v_j\} \in E} (1 - x_j)),$$

where this sum brings  $P_G(\vec{x})$  to infinity whenever there is a non-dominated vertex which is not in the selected dominating set. In addition,  $P_G(\vec{x}) \leq k$  then guarantees a dominating set of size at most  $k$ .

- (4) Similar observations as for VERTEX COVER, INDEPENDENT SET, and DOMINATING SET do hold for many other graph problems and, in particular, weighted variants of these.<sup>15</sup> As a source of problems, consider the variants of DOMINATING SET listed in [46]. In particular, the TOTAL DOMINATING SET problem is defined by

$$P_G^{\text{ver}}(\vec{x}) = \sum_{v_i \in V} (\infty \cdot \prod_{\{v_i, v_j\} \in E} (1 - x_j)).$$

Moreover, graph problems where a small (or large) *edge set* is sought for can often be reformulated into vertex set optimization problems by introducing an additional artificial vertex on each edge of the original graph. In this way, the EDGE DOMINATING SET [47] problem can be handled. Similarly, planar graph problems where a small (or large) *face set* is looked for are expressible as select&verify problems of the dual graphs.

We will also need a notion of select&verify problems where the “selecting function” and the “verifying function” operate on a subgraph of the given graph.

**Definition 20** Let  $P = P^{\text{sel}} + P^{\text{ver}}$  be the function of a select&verify problem. For an  $n$ -vertex graph  $G$  and subgraphs  $G^{\text{ver}} = (V^{\text{ver}}, E^{\text{ver}})$ ,  $G^{\text{sel}} = (V^{\text{sel}}, E^{\text{sel}}) \subseteq G$ , we let

$$P_{G^{\text{ver}}}(\vec{x} \mid G^{\text{sel}}) := P_{G^{\text{ver}}}^{\text{ver}}(\pi_{V^{\text{ver}}}(\vec{x})) + P_{G^{\text{sel}}}^{\text{sel}}(\pi_{V^{\text{sel}}}(\vec{x})),$$

where  $\pi_{V'}$  is the projection of the vector  $\vec{x} \in \{0, 1\}^n$  to the variables corresponding to the vertices in  $V'$ .

<sup>15</sup>In the weighted case, one typically chooses a “selecting function” of the form  $P_G^{\text{sel}} = \sum_{v_i \in V} \alpha_i x_i$ , where  $\alpha_i$  is the weight of the vertex  $v_i$ .

### Weakly glueable graph problems

We will solve graph problems, slicing the given graph into small pieces with the help of small separators. Within these separators, the basic strategy will be to test all possible assignments of the vertices. The separators will serve as boundaries between the different graph chunks into which the graph is split. For each possible assignment of the vertices in the separators, we want to—independently—solve the corresponding problems on the graph chunks and then reconstruct a solution for the whole graph by “gluing” together the solutions for the graph chunks. Therefore, all additional information necessary for solving the subproblems correctly has to be transported and coded within the separators. It turns out that the information to be passed on is pretty clear in the case of the VERTEX COVER problem, but it is much more involved in the case of the DOMINATING SET problem and many others. This is the basic motivation for the formal framework we develop here. We need to assign *colors* to the separator vertices in the course of the algorithm. Hence, our algorithm has to be designed in such a manner that it can also cope with colored graphs, even though the original problem may have been a problem on non-colored graphs. In general (e.g., in the case of the DOMINATING SET problem), it is not sufficient to simply use the two colors 1 (for encoding “in the selected set”) and 0 (for “not in the selected set”).

**Definition 21** *Let  $G = (V, E)$  be an undirected graph and let  $C_0, C_1$  be finite, disjoint sets. A  $C_0$ - $C_1$ -coloring of  $G$  is a function  $\chi : V \rightarrow C_0 + C_1 + \{\#\}$ .<sup>16</sup> A  $C_0$ - $C_1$ -coloring with  $C_0 = \{0\}$  and  $C_1 = \{1\}$  is called a 0-1-coloring.*

*For a  $C_0$ - $C_1$ -coloring  $\chi$ , the corresponding 0-1-coloring  $\hat{\chi}$  is given by*

$$\hat{\chi}(v) = \begin{cases} i & \text{if } \chi(i) \in C_i, \quad i = 0, 1, \\ \# & \text{otherwise.} \end{cases}$$

For  $V' \subseteq V$ , a function  $\chi : V' \rightarrow C_0 + C_1$  can naturally be extended to a  $C_0$ - $C_1$ -coloring of  $G$  by setting  $\chi(v) = \#$  for all  $v \in V \setminus V'$ .

**Definition 22** *For two 0-1-colorings  $\chi_1, \chi_2 : V \rightarrow \{0, 1, \#\}$  with  $\chi_1^{-1}(\{0, 1\}) \cap \chi_2^{-1}(\{0, 1\}) = \emptyset$ , the sum  $\chi_1 + \chi_2$  is defined by*

$$(\chi_1 + \chi_2)(v) = \begin{cases} \chi_1(v) & \text{if } \chi_1(v) \neq \#, \\ \chi_2(v) & \text{if } \chi_2(v) \neq \#, \\ \# & \text{otherwise.} \end{cases}$$

**Definition 23** *Consider an instance  $(G, k)$  of a select&verify problem  $\mathcal{G}$  and*

<sup>16</sup>The symbol  $\#$  will be used for the undefined (i.e., not yet defined) color.

a vector  $\vec{x} \in \{0, 1\}^n$  with  $n = |V(G)|$ . Let  $\chi$  be a 0-1-coloring of  $G$ . Then,  $\vec{x}$  is consistent with  $\chi$ , written  $\vec{x} \sim \chi$ , if

$$\chi(v_j) = i \Rightarrow x_j = i, \quad \text{for } i = 0, 1, j = 1, \dots, n.$$

We now provide the central notion of “weakly glueable” select&verify problems, a weaker form of the glueability concept as introduced in [6]. We apply this rather abstract notion to concrete graph problems afterwards.

**Definition 24** A select&verify problem  $\mathcal{G}$  given by  $(P, \text{opt})$  is weakly glueable with  $\lambda$  colors if there exist

- a color set  $C := C_0 + C_1 + \{\#\}$  with  $|C_0 + C_1| = \lambda$ , and
- a polynomial time computable function  $h : (\mathbb{R}_+ \cup \{\pm\infty\})^3 \rightarrow \mathbb{R}_+ \cup \{\pm\infty\}$ ;

and, for every  $n$ -vertex graph  $G = (V, E)$  and subgraphs  $G^{sel}, G^{ver} \subseteq G$  with a separation  $(A_1, S, A_2)$  of  $G^{ver}$ , we find, for each coloring  $\chi : S \rightarrow C_0 + C_1$ ,

- subgraphs  $G(A_i, \chi)$  of  $G^{ver}$  with  $G^{ver}[A_i] \subseteq G(A_i, \chi) \subseteq G^{ver}[\delta A_i]$  for  $i = 1, 2$ ,
- subgraphs  $G(S, \chi)$  of  $G^{ver}$  with  $G(S, \chi) \subseteq G^{ver}[S]$

such that, for each 0-1-coloring  $\mu : V \rightarrow \{0, 1, \#\}$  with  $\mu|_S \equiv \#$ , we have

$$\text{opt}_{\substack{\vec{x} \in \{0,1\}^n \\ \vec{x} \sim \mu}} P_{G^{ver}}(\vec{x} \mid G^{sel}) = \text{opt}_{\chi: S \rightarrow C_0 + C_1} h\left(\text{Eval}_{A_1}(\chi), \text{Eval}_S(\chi), \text{Eval}_{A_2}(\chi)\right). \quad (3)$$

Here,  $\text{Eval}_X(\cdot)$  for  $X \in \{A_1, S, A_2\}$  is of the form

$$\text{Eval}_X(\chi) = \text{opt}_{\substack{\vec{x} \in \{0,1\}^n \\ \vec{x} \sim (\mu + \chi)}} P_{G(X, \chi)}(\vec{x} \mid G[X] \cap G^{sel}).$$

**Example 25** We give some examples of weakly glueable problems, where — for the time being — we restrict ourselves to the case where  $G^{ver} = G^{sel} = G$ . The subtlety of allowing different subgraphs  $G^{ver}, G^{sel}$  in the definition above is due to technical reasons that become clear later. All examples generalize in a straight-forward way to this case.

- (1) VERTEX COVER is weakly glueable with  $\lambda = 2$  colors. We use the color sets  $C_i := \{i\}$  for  $i = 0, 1$ . The function  $h$  is  $h(x, y, z) = x + y + z$ . The subgraphs  $G(X, \chi)$  for  $X \in \{A_1, S, A_2\}$  and  $\chi : S \rightarrow C_0 + C_1$  are

$$\begin{aligned} G(A_i, \chi) &:= G[A_i \cup \chi^{-1}(0)] \quad \text{for } i = 1, 2, \quad \text{and} \\ G(S, \chi) &:= G[S]. \end{aligned}$$

The subroutine  $\text{Eval}_S(\chi)$  checks if the coloring  $\chi$  yields a vertex cover on  $G[S]$  and the subroutines  $\text{Eval}_{A_i}(\chi)$  compute the minimum size vertex

cover on  $G[A_i]$  under the constraint that all neighbors in  $A_i$  of a vertex in  $\chi^{-1}(0)$  are covered. Obviously, Eq. (3) is thus satisfied.

- (2) Similarly, INDEPENDENT SET is weakly glueable with 2 colors.  
 (3) DOMINATING SET is weakly glueable with  $\lambda = 4$  colors, using  $C_0 := \{0_{A_1}, 0_{A_2}, 0_S\}$  and  $C_1 := \{1\}$ . The semantics of these colors is as follows. Assigning the color  $0_X$ , for  $X \in \{A_1, S, A_2\}$ , to vertices in a separation  $(A_1, S, A_2)$  means that the vertex is not in the dominating set and will be dominated by a vertex from  $X$ . Color 1 means that the vertex belongs to the dominating set. We set  $h(x, y, z) = x + y + z$ . For a given coloring  $\chi : S \rightarrow C_0 + C_1$ , we define

$$G(A_i, \chi) := G[A_i \cup \chi^{-1}(\{1, 0_{A_i}\})]$$

$$G(S, \chi) := G[\chi^{-1}(\{1, 0_S\})].$$

In this way, color information is passed to the subproblems.  $Eval_S(\chi)$  checks whether the assignments of the color  $0_S$  are correct (i.e., whether all vertices assigned  $0_S$  are dominated by a vertex from  $S$ ). Also,  $Eval_{A_i}$  returns the size of a minimum dominating set in  $A_i$  under the constraint that some vertices in  $\delta A_i$  still need to be dominated (namely, the vertices in  $\chi^{-1}(0_{A_i})$ ) and some vertices in  $\delta A_i$  can already be assumed to be in the dominating set (namely, the vertices in  $\chi^{-1}(1)$ ). With these settings, Eq. (3) is satisfied.

We want to mention in passing that—besides the problems given here—many more select&verify problems are weakly glueable. In particular this is true for the weighted versions and variations of the above mentioned problems. Note that, e.g., TOTAL DOMINATING SET is an example of a graph problem where a color set  $C_1$  of more than one color is needed.

### 5.2.2 The algorithm

Similar to Theorem 16, which is based on tree decompositions, we construct a partial layerwise separation  $\mathcal{S}(\psi)$  with optimally adapted trade-off parameter  $\psi$  to guarantee an efficient dynamic programming algorithm. However, for our purposes here, we need to be able to deal with “precolored” graphs.

**Definition 26** Let  $\mathcal{G}$  be a select&verify graph problem defined by  $(P, \text{opt})$ . The problem CONSTRAINT  $\mathcal{G}$  then is to determine, for an  $n$ -vertex graph  $G = (V, E)$ , two subgraphs  $G^{sel}, G^{ver} \subseteq G$ , and a given 0-1-coloring  $\chi : V \rightarrow \{0, 1, \#\}$ , the value

$$\text{opt}_{\substack{\vec{x} \in \{0,1\}^n \\ \vec{x} \sim \chi}} P_{G^{ver}}(\vec{x} \mid G^{sel}).$$

**Theorem 27** Let  $\mathcal{G}$  be a select&verify problem for planar graphs. Suppose that




Fig. 2. Dynamic programming on layerwisely separated graph.

- (1)  $\mathcal{G}$  has the LSP of width  $w$  and size-factor  $d$ ,
- (2)  $\mathcal{G}$  is weakly glueable with  $\lambda$  colors, and
- (3) there exists an algorithm that solves the problem CONSTRAINT  $\mathcal{G}$  for a given graph  $G$  in time  $\tau^{\text{out}(G)}n$ .

Then, there is an algorithm to decide  $(G, k) \in \mathcal{G}$  in time  $O(\tau^w \cdot 2^{\theta_2(\lambda, \tau, d)\sqrt{k}}n)$ , where  $\theta_2(\lambda, \tau, d) = 2\sqrt{d \log(\lambda) \log(\tau)}$ .

**PROOF.** Let us first sketch the overall structure of the algorithm.

- (1) Compute some planar embedding  $\phi$  of  $G$ , and find a “suitable” partial layerwise separation  $(S_1, \dots, S_q)$  for the plane graph  $(G, \phi)$ . A coarse sketch of the such obtained graph structure is depicted in Figure 2.
- (2) By using dynamic programming techniques, an optimal solution is found by sweeping over the graph from left to right, as illustrated in Figure 2. More detailed, we do the following:
  - (a) For all possible “colorings” of  $S_1$ , find an optimal solution of CONSTRAINT  $\mathcal{G}$  on  $G_0$  (plus suitably chosen precolored vertices from  $S_1$ ); store the obtained optima in a (large) table belonging to  $S_1$ .
  - (b) For  $j := 2$  to  $q$  do:
    - For all possible “colorings” of  $S_{j-1}$  and of  $S_j$ , find an optimal solution of CONSTRAINT  $\mathcal{G}$  on  $G_{j-1}$  (plus suitably chosen precolored vertices from  $S_{j-1}$  as well as of  $S_j$ ).
    - Store the obtained optima (for the subgraph of  $G$  with vertices from  $G_0$  through  $G_{j-1}$  and  $S_1$  through  $S_j$ ) in a table belonging to  $S_j$ .
    - (For reasons of space efficiency, you might now forget about the table belonging to  $S_{j-1}$ .)
  - (c) For all possible “colorings” of  $S_q$ , find an optimal solution of CONSTRAINT  $\mathcal{G}$  on  $G_q$  (plus suitably chosen precolored vertices from  $S_q$ ); store the obtained optima in a (large) table belonging to  $S_q$ .
  - (d) From the table pertaining to  $S_q$ , the desired optimum is found.

We are now giving formal details of the sketched algorithms, thereby proving its correctness. Suppose  $\mathcal{G}$  is defined by  $(P, \text{opt})$ .

**Step 1:** Given an instance  $(G, k)$ , in linear time we can compute some planar embedding  $\phi$  of  $G$  (see [18]). Compute a partial layerwise separation  $\mathcal{S}(\psi)$  ( $\psi$  will be determined later) for  $(G, \phi)$ , using Proposition 13 and the assumption 1 (LSP). Suppose  $\mathcal{S}(\psi) = (S_1, \dots, S_q)$  and let  $\mathcal{C}_{\mathcal{S}(\psi)} = (G_0, \dots, G_q)$  denote the corresponding graph-chunks cut out by  $\mathcal{S}(\psi)$ .

For every separator  $S_i$ , we get a separation of the form

$$(A_{\text{out}}^{(i)}, S_i, A_{\text{in}}^{(i)}),$$

where  $A_{\text{out}}^{(i)}$ , and  $A_{\text{in}}^{(i)}$ , respectively, are the vertices of the graph chunks of lower order layers, and higher order layers, respectively. By default, we let  $S_0 = S_{q+1} = \emptyset$  such that the corresponding separations are  $(\emptyset, S_0, V)$  and  $(V, S_{q+1}, \emptyset)$ , respectively.

The layerwise separation and the separations  $(A_{\text{out}}^{(i)}, S_i, A_{\text{in}}^{(i)})$  are illustrated in Figure 2.

**Step 2:** Determine the value

$$\text{opt}_{\vec{x} \in \{0,1\}^n} P_G(\vec{x}).$$

This can be done by a dynamic programming approach that makes use of the weak glueability of  $\mathcal{G}$  as follows.

We successively compute, for  $i = 1, \dots, q + 1$ , the values

$$M^{(i)}(\mu^{(i)}) := \text{opt}_{\vec{x} \sim \widehat{\mu^{(i)}}} P_{H_i^{\text{ver}}(\mu^{(i)})}(\vec{x} \mid H_i^{\text{sel}}) \quad (4)$$

for all  $C_0$ - $C_1$ -colorings  $\mu^{(i)} : S_i \rightarrow C_0 + C_1$ , where

$$H_i^{\text{ver}}(\mu^{(i)}) := G(A_{\text{out}}^{(i)}, \mu^{(i)}) \quad \text{and} \quad H_i^{\text{sel}} := G[A_{\text{out}}^{(i)}].$$

Note that we have

$$\text{opt}_{\vec{x} \in \{0,1\}^n} P_G(\vec{x}) = M^{(q+1)}(\mu),$$

for the empty map  $\mu : S_{q+1} = \emptyset \rightarrow C_0 + C_1$ , because  $H_{q+1}^{\text{sel}} = G[A_{\text{out}}^{(q+1)}] = G[V] = G$  and  $H_{q+1}^{\text{ver}}(\mu) = G$  (since  $H_{q+1}^{\text{sel}} \subseteq H_{q+1}^{\text{ver}}(\mu)$ ).

The computation of  $M^{(i)}(\mu^{(i)})$  as defined in (4) can be done iteratively. To do so, note that  $H_i^{\text{ver}}(\mu^{(i)})$  is separated by  $S_{i-1}$  in

$$(B_{\text{out}}^{(i-1)}, S_{i-1}, B_{\text{in}}^{(i-1)}),$$



where  $B_{\text{out}}^{(i-1)} = A_{\text{out}}^{(i-1)}$  and  $B_{\text{in}}^{(i-1)} = A_{\text{in}}^{(i-1)} \cap V(H_i^{\text{ver}}(\mu^{(i)}))$ . Hence, by definition of weak glueability we have

$$M^{(i)}(\mu^{(i)}) = \text{opt}_{\mu^{(i-1)}:S_{i-1} \rightarrow C_0+C_1} h(\text{Eval}_{B_{\text{out}}^{(i-1)}}(\mu^{(i-1)}), \text{Eval}_{S_{i-1}}(\mu^{(i-1)}), \text{Eval}_{B_{\text{in}}^{(i-1)}}(\mu^{(i-1)})), \quad (5)$$

where

$$\text{Eval}_X(\mu^{(i-1)}) = \text{opt}_{\vec{x} \sim (\widehat{\mu^{(i-1)}} + \widehat{\mu^{(i)}})} P_{G(X, \mu^{(i-1)})}(\vec{x} \mid G[X] \cap H_i^{\text{sel}}). \quad (6)$$

for  $X \in \{B_{\text{out}}^{(i-1)}, S_{i-1}, B_{\text{in}}^{(i-1)}\}$ . Here, recall that  $\widehat{\mu^{(i)}}$  denotes the 0-1-coloring corresponding to  $\mu^{(i)}$ . In particular, for the different choices of  $X$ , we get the following.

- For  $X = B_{\text{out}}^{(i-1)}$ , equation (6) becomes

$$\begin{aligned} \text{Eval}_{B_{\text{out}}^{(i-1)}}(\mu^{(i-1)}) &= \text{opt}_{\vec{x} \sim (\widehat{\mu^{(i-1)}} + \widehat{\mu^{(i)}})} P_{H_{i-1}^{\text{ver}}(\mu^{(i-1)})}(\vec{x} \mid H_{i-1}^{\text{sel}} \cap H_i^{\text{sel}}) \\ &= M^{(i-1)}(\mu^{(i-1)}), \end{aligned} \quad (7)$$

where the last equation holds, since  $H_{i-1}^{\text{sel}} \subseteq H_i^{\text{sel}}$  and since  $\widehat{\mu^{(i)}} \equiv \#$  restricted to  $G(B_{\text{out}}^{(i-1)}, \mu^{(i-1)}) = H_{i-1}^{\text{ver}}(\mu^{(i-1)})$ .

Hence, the value of  $\text{Eval}_{B_{\text{out}}^{(i-1)}}(\mu^{(i-1)})$  is given by the previous step of the iteration.

- For  $X = S_{i-1}$ , equation (6) becomes

$$\begin{aligned} \text{Eval}_{S_{i-1}}(\mu^{(i-1)}) &= \text{opt}_{\vec{x} \sim \widehat{\mu^{(i-1)}}} P_{G(S_{i-1}, \mu^{(i-1)})}(\vec{x} \mid G[S_{i-1}]) \\ &= P_{G(S_{i-1}, \mu^{(i-1)})}(\vec{x}_{\widehat{\mu^{(i-1)}}} \mid G[S_{i-1}]), \end{aligned} \quad (8)$$

where  $\vec{x}_{\widehat{\mu^{(i-1)}}} \in \{0, 1\}^n$  is an arbitrary vector such that  $\vec{x}_{\widehat{\mu^{(i-1)}}} \sim \widehat{\mu^{(i-1)}}$ . The first equation above holds, since  $G(S_{i-1}, \mu^{(i-1)}) \subseteq G[S_{i-1}] \subseteq H_i^{\text{sel}}$  and  $\widehat{\mu^{(i)}} \equiv \#$  restricted to  $G(S_{i-1}, \mu^{(i-1)})$ . The second equation is true since the 0-1-coloring  $\widehat{\mu^{(i-1)}}$  assigns color 0 or color 1 to all vertices in  $S_{i-1}$ , and since  $G(S_{i-1}, \mu^{(i-1)}) \subseteq G[S_{i-1}]$ .

Hence, the value  $\text{Eval}_{S_{i-1}}(\mu^{(i-1)})$  can be computed by a simple evaluation of the function  $P$  for the given vector  $\vec{x}_{\widehat{\mu^{(i-1)}}}$ .

- For  $X = B_{\text{in}}^{(i-1)}$ , equation (6) becomes

$$\begin{aligned} \text{Eval}_{B_{\text{in}}^{(i-1)}}(\mu^{(i-1)}) &= \text{opt}_{\vec{x} \sim (\widehat{\mu^{(i-1)}} + \widehat{\mu^{(i)}})} P_{G(B_{\text{in}}^{(i-1)}, \mu^{(i-1)})}(\vec{x} \mid G[B_{\text{in}}^{(i-1)}] \cap H_i^{\text{sel}}) \\ &= \text{opt}_{\vec{x} \sim (\widehat{\mu^{(i-1)}} + \widehat{\mu^{(i)}})} P_{G(\mu^{(i)}, \mu^{(i-1)})}(\vec{x} \mid G_{i-1}), \end{aligned} \quad (9)$$




Fig. 3. The graph  $G(\mu^{(i)}, \mu^{(i-1)})$ .

where  $G(\mu^{(i)}, \mu^{(i-1)}) := G(A_{\text{in}}^{(i-1)}, \mu^{(i-1)}) \cap G(A_{\text{out}}^{(i)}, \mu^{(i)})$ . This graph is illustrated in Figure 3. In the evaluation above we used that  $G[B_{\text{in}}^{(i-1)}] \cap H_i^{\text{sel}} = G_{i-1}$  and that  $G(B_{\text{in}}^{(i-1)}, \mu^{(i-1)}) = G(\mu^{(i)}, \mu^{(i-1)})$ .

Hence, the value  $\text{Eval}_{B_{\text{in}}^{(i-1)}}(\mu^{(i-1)})$  can be computed using the  $\tau^{\text{out}(G)}$  time algorithm for CONSTRAINT  $\mathcal{G}$ .

Hence, plugging formulas (7), (8), and (9) in expression (5), we obtain

$$M^{(i)}(\mu^{(i)}) = \text{opt}_{\mu^{(i-1)}: S_{i-1} \rightarrow C_0 + C_1} h \left( \begin{array}{c} M^{(i-1)}(\mu^{(i-1)}) \\ P_{G(S_{i-1}, \mu^{(i-1)})}(\vec{x}_{\widehat{\mu^{(i-1)}}} \mid G[S_{i-1}]) \\ \text{opt}_{\vec{x} \sim \widehat{\mu^{(i-1)}} + \widehat{\mu^{(i)}}} P_{G(\mu^{(i)}, \mu^{(i-1)})}(\vec{x} \mid G_{i-1}) \end{array} \right). \quad (10)$$

This evaluation is done successively for all  $i = 1, \dots, q + 1$ . By induction, one sees that

$$\text{opt}_{\vec{x} \in \{0,1\}^n} P_G(\vec{x}) = M^{(q+1)}(\mu)$$

can be computed in this way.

**Computation time:** For fixed coloring  $\mu^{(i)}$ , computing  $M^{(i)}(\mu^{(i)})$  according to equation (10) costs time

$$\lambda^{|S_{i-1}|} \cdot \tau^{\text{out}(G[S_{i-1} \cup V_{G_{i-1}} \cup S_i])}.$$

The first factor reflects the cost of computing the opt over all  $\mu^{(i-1)} : S_{i-1} \rightarrow C_0 + C_1$ . The second factor arises by the evaluations on the graphs  $G(S_{i-1}, \mu^{(i-1)})$ ,  $G(\mu^{(i-1)}, \mu^{(i)}) \subseteq G[S_{i-1} \cup V_{G_{i-1}} \cup S_i]$ , where we use assumption 3 of the theorem. Thus, the running time for evaluating  $M^{(i)}(\mu^{(i)})$  for all  $\mu^{(i)} : S_i \rightarrow C_0 + C_1$  is bounded by

$$\lambda^{|S_i|} \cdot \lambda^{|S_{i-1}|} \cdot \tau^{\text{out}(G[S_{i-1} \cup V_{G_{i-1}} \cup S_i])}.$$

Hence, the total running time of the algorithm is bounded by  $2^{\theta(\psi)} n$ , where

$$\begin{aligned}
 \theta(\psi) &\leq \log(\lambda) \sum_{i=1, \dots, q} |S_i| + \log(\tau) \max_{i=0, \dots, q} \text{out}(G[S_i \cup V_{G_i} \cup S_{i+1}]) \\
 &\leq \log(\lambda) \psi \sqrt{dk} + \log(\tau) \left( \frac{\sqrt{dk}}{\psi} + w \right) \\
 &= \left( \log(\lambda) \psi + \frac{\log \tau}{\psi} \right) \sqrt{dk} + \log(\tau) w
 \end{aligned}$$

This upper bound is minimized for  $\psi = \sqrt{\log(\tau)/\log(\lambda)}$ , which gives us the claimed value  $\theta_2(\lambda, \tau, d) = 2\sqrt{d \log(\lambda) \log(\tau)}$  and the constant  $\tau^w$ .

It remains to say for which problems there exists a solving algorithm of the problem CONSTRAINT  $\mathcal{G}$  for a given graph  $G$  in time  $\tau^{\text{out}(G)} n$ . Baker [10] presented several such problems  $\mathcal{G}$ , however, the algorithms as stated there do *not* cope with the constraint version of  $\mathcal{G}$ . Often, we can adapt them quite easily. In the case of PVS, as well as in the case of PIS, it is quite simple to handle a “precoloring”  $\mu : V \rightarrow \{0, 1, \#\}$  for a graph  $G = (V, E)$ .

More formally, given an admissible<sup>17</sup> coloring  $\mu$ , one likes to transform an instance  $(G, k)$  to an instance  $(G', k')$ , such that  $(G, k) \in \mathcal{G}$  for some witnessing vector  $\vec{x}$  with  $\vec{x} \sim \mu$  iff  $(G', k') \in \mathcal{G}$  for some witnessing vector  $\vec{x}'$  (without any constraint).

For PVS, this can, e.g., be achieved by the following transformation:

$$\begin{aligned}
 G' &= G[V - (C \cup N(\mu^{-1}(0)))] \quad \text{and} \\
 k' &= k - |\mu^{-1}(1)| - |\{v \in V - C \mid \exists u \in \mu^{-1}(0) \cap N(v)\}|,
 \end{aligned}$$

here  $C := \mu^{-1}(\{0, 1\})$  denotes the vertices that are already assigned a color. A vertex  $v \in C$ , which is—by the coloring  $\mu$ —assigned not to be in the vertex cover, i.e.,  $\mu(v) = 0$ , forces its neighbors to be in a vertex cover. Hence, the set  $N(\mu^{-1}(0))$  needs to be in any vertex cover (given by  $\vec{x}$ ) that fulfills the constraint  $\vec{x} \sim \mu$ . This justifies the definition of  $G'$ . The parameter  $k$  becomes smaller by the number of vertices which are already assigned to be in the vertex cover, i.e.,  $\mu^{-1}(1)$ , and the number of vertices that are forced to be in the vertex cover by  $\mu$ , i.e., by the number of vertices in  $V - C$  that have a neighbor in  $\mu^{-1}(0)$ . We can apply the non-constraint algorithm to  $(G', k')$ , with  $\text{out}(G') \leq \text{out}(G)$ . A similar observation helps deal with PIS.

**Example 28** (1) For PVS, we have  $d = 2$ ,  $w = 1$  (see Example 11.1),  $\lambda = 2$  (see Example 25.1), and  $\tau = 8$  (see the result of Baker [10] which can be

<sup>17</sup> Here, *admissible* means that there exists a vector  $\vec{x} \in \{0, 1\}^n$  with  $\vec{x} \sim \mu$ , such that  $P_G^{\text{ver}}(\vec{x}) = 0$ .

adapted to the constraint case by the considerations above) and, hence, the approach in Theorem 27 yields an  $O(2^{2\sqrt{6k}}n)$  time algorithm.

- (2) Similarly, Examples 11.2, and 25.2 give  $d = 4$ ,  $w = 1$  and  $\lambda = 2$  for PIS. Since  $\tau = 8$  (see [10] and our considerations for the constraint case), we obtain a  $O(2^{4\sqrt{3k}}n)$  time algorithm.
- (3) Kanj and Perkiovič have recently shown [31] that CONSTRAINT DOMINATING SET can be solved in  $O(27^{\text{out}(G)}n)$  time, this way improving on Example 17. Together with Example 9.1, Theorem 27 gives an  $O(3^{6\sqrt{17k}}n) \approx O(2^{39.21\sqrt{k}}n)$  time algorithm for PDS.

### 5.3 Comparing the benefits of the different approaches

Which of the two approaches (presented in Subsections 5.1 and 5.2, respectively) for the algorithms on layerwisely separated graphs should be preferred? For that purpose, let us compare the results obtained in Theorems 16 and 27.

Clearly, both results rely on the LSP assumption. Besides that, Theorem 16 requires the existence of a linear time algorithm for bounded treewidth graphs, whereas in Theorem 27 one needs (besides the weak glueability assumption) the existence of a linear time algorithm for graphs with bounded outerplanarity.<sup>18</sup> The interrelation inbetween these assumptions is given by the following observation, which is based on Proposition 14.

**Lemma 29** *Let  $\mathcal{G}$  be a parameterized problem for planar graphs. Suppose that there exists a time  $\sigma^\ell n$  algorithm that solves CONSTRAINT  $\mathcal{G}$ , when graph  $G$  is given together with a tree decomposition of width  $\ell$ . Then, there is an algorithm that solves CONSTRAINT  $\mathcal{G}$  in time  $\tau^{\text{out}(G)}n$  for  $\tau = \sigma^3$ .*

**PROOF.** Applying the algorithm of Proposition 14 to input instance  $(G, k)$ , we obtain a tree decomposition of width at most  $3 \text{out}(G)$ . Then CONSTRAINT  $\mathcal{G}$  can be solved using this tree decomposition by our assumption.

The following easy corollary helps compare the approach from Subsection 5.1 (i.e., Theorem 16) with the approach in Subsection 5.2 (i.e., Theorem 27).

---

<sup>18</sup> Clearly, for both results, Theorem 16 and Theorem 27, respectively, we do not rely on a *linear* time algorithm for bounded treewidth graphs, and graphs with bounded outerplanarity, respectively. Assuming a polynomial time algorithms is sufficient to get a  $c^{\sqrt{k}}$ -algorithm. The results are formulated for linear time algorithm since we do have linear time algorithms here.

**Corollary 30** *Let  $\mathcal{G}$  be a select&verify problem for planar graphs. Suppose that*

- (1)  $\mathcal{G}$  has the LSP of width  $w$  and size-factor  $d$ ,
- (2)  $\mathcal{G}$  is weakly glueable with  $\lambda$  colors, and
- (3) there exists a time  $\sigma^\ell n$  algorithm that solves CONSTRAINT  $\mathcal{G}$  for a graph  $G$ , if  $G$  is given together with a tree decomposition of width  $\ell$ .

Then, there is an algorithm to decide  $(G, k) \in \mathcal{G}$  in time  $O(\sigma^{3w} \cdot 2^{\theta_3(\lambda, \sigma, d)\sqrt{k}n})$ , where  $\theta_3(\lambda, \sigma, d) = 2\sqrt{3d \log(\lambda) \log(\sigma)}$ .

The exponential factor of the algorithm in Corollary 30, i.e.,  $\theta_3(\lambda, \sigma, d)$ , is related to the corresponding exponent of Theorem 16, i.e.,  $\theta_1(\sigma, d)$  in the following way:

$$\sqrt{\log \lambda} \cdot \theta_1(\sigma, d) = \sqrt{\log \sigma} \cdot \theta_3(\lambda, \sigma, d).$$

From this we derive that

- if  $\lambda > \sigma$ , the algorithm in Theorem 16 outperforms the result of Corollary 30,
- if  $\lambda < \sigma$ , the algorithm in Corollary 30 outperforms the result of Theorem 16.

However, in order to apply Corollary 30, we need the extra assumptions that we have a select&verify problem which is weakly glueable and that we can deal with the problem CONSTRAINT  $\mathcal{G}$  in the treewidth algorithm.

For our running examples PVS, PIS and PDS, it is possible to apply Corollary 30 instead of Theorem 27. We suppress details here, since both strategies lead to the same constants (as derived in Example 28). Observe that, for algorithmic purposes, three colors are enough in the case of PDS.

We now remark on the *space consumption* of the two different approaches of Theorem 16, and Theorem 27, respectively. The tables that need to be maintained in order to realize a treewidth based algorithm of Theorem 16 (i.e., the tables corresponding to the bags of the underlying tree decomposition) have size bounded by

$$\sigma^{2\sqrt{6dk} + (3w-1)}.$$

In contrast, the analysis of the proof of Theorem 27 gives an upper bound on the size of the tables for the dynamic programming therein. The upper bound is given by

$$\max_{i=1, \dots, q} \lambda^{|S_i|} \leq \lambda \sqrt{\frac{\log(\tau)dk}{2 \log(\lambda)}}.$$

Since all our examples have the property that  $\sigma = \lambda$ , in terms of space consumption, the approach of Theorem 27 outperforms Theorem 16.

**Extending our results to graph classes containing non-planar graphs** (as done in [19]) necessitates the employment of treewidth based techniques,

since the “geometrical” notion of outerplanarity is bound to planar graphs.

#### 5.4 Beyond linear kernels

This section sketches an extension of the methods presented so far.

As already mentioned after Definition 18, every select&verify problem  $\mathcal{G}$  that admits a linear problem kernel is solvable in time  $2^{O(k)}n^{O(1)}$ . Also, the existence of a linear problem kernel implies the LSP. Assuming certain further properties of  $\mathcal{G}$ , we have seen results which improve the running time of a solving algorithm to  $2^{O(\sqrt{k})}n^{O(1)}$ .

The techniques can be extended to the case where  $\mathcal{G}$  admits a problem kernel of size  $p(k)$ . We briefly outline how, by a straightforward generalization of our methods, the running time for solving  $\mathcal{G}$  can be sped up from  $2^{O(p(k))}n^{O(1)}$  (as can be obtained trivially) to  $2^{O(\sqrt{p(k)})}n^{O(1)}$ . The key tool for that purpose is the extension of Proposition 13, which now becomes:

**Proposition 31** *Let  $(G = (V, E), \phi)$  be a plane graph that admits a layerwise separation of width  $w$  and size  $p(k)$ . Then, for every  $\psi \in \mathbb{R}_+$ , there exists a partial layerwise separation  $\mathcal{S}(\psi)$  of width  $w$  such that*

- (1)  $\sum_{S \in \mathcal{S}(\psi)} |S| \leq \psi \sqrt{p(k)}$  and
- (2)  $\text{out}(H) \leq \frac{\sqrt{p(k)}}{\psi} + w$  for each graph chunk  $H$  in  $\mathcal{C}_{\mathcal{S}(\psi)}$ .

Moreover, there is an algorithm with running time  $O(\sqrt{p(k)}n)$  which, for a given  $\psi$ ,

- recognizes whether  $(G, \phi)$  admits a layerwise separation of width  $w$  and size  $p(k)$  and, if so, computes  $\mathcal{S}(\psi)$ ;
- computes a partial layerwise separation of width  $w$  that fulfills the conditions above.

As in Section 5, the above result can be used to show similar results as those given in Subsection 5.1 (see Theorem 16) and in Subsection 5.2 (see Theorem 27).

Note that in both theorems it would be sufficient to replace condition 1 by a modified version of the LSP:

- 1'. for every  $(G, k) \in \mathcal{G}$  the graph  $G$  has a layerwise separation of width  $w$  and size  $p(k)$ .

Using similar arguments as in the proof of Lemma 10, having a problem kernel of size  $p(k)$  implies condition 1'. on the set of all problem kernels  $\mathcal{G}'$ . With this modification—omitting details— $O(c\sqrt{p(k)}|G|)$  algorithms can be achieved.

## 6 Conclusion

This paper presents new, improved results for the fixed-parameter complexity of planar graph problems. To some extent, this paper can be seen as the “parameterized complexity counterpart” to what was developed by Baker [10] in the context of approximation algorithms. We describe two main ways (namely linear problem kernels and problem-specific approaches) to achieve the novel concept of Layerwise Separation Property, from which again, two approaches (tree decomposition and bounded outerplanarity) lead to  $c^{\sqrt{k}}$ -algorithms for planar graph problems (see Figure 1 for an overview). A slight modification of our presented techniques can be used to extend our results to parameterized problems that admit a problem kernel of size  $p(k)$ . Then, the running time can be sped up from  $2^{O(p(k))}n^{O(1)}$  to  $2^{O(\sqrt{p(k)})}n^{O(1)}$ . It appears to be that all FPT-problems that admit treewidth based algorithms [12,13,45,46] can be handled by our methods.

Future research topics raised by our work include, among others,

- to further improve the (“exponential”) constants, e.g., by a further refined and more sophisticated “layer decomposition tree”; as already mentioned in the introduction, lots of work in this direction has been put on its way since the conference version of this paper appeared; more specifically, for PDS and PVC, the branchwidth approach seems to yield slightly better constants (by a factor of  $\sqrt{3}/1.5$  in the exponent) than the treewidth/outerplanarity approach;<sup>19</sup> and
- to investigate and extend the availability of linear problem kernels for all kinds of planar graph problems.

Finally, a more general question is whether there are other “problem classes” that allow for  $c^{\sqrt{k}}$  fixed-parameter algorithms. Cai and Juedes [16] addressed this issue and claimed that for a list of parameterized problems (e.g., VERTEX COVER on general graphs)  $c^{o(k)}$ -algorithms are impossible unless  $\text{FPT} = W[1]$ . However, a flaw was detected in their proof [23] and it has been corrected in [15]. Now the non-existence of  $c^{\sqrt{k}}$ -algorithms is based on the conjecture

---

<sup>19</sup>The claimed bound for PDS in the treewidth/outerplanarity approach is due to Fernau and Juedes (ongoing work), making use of the more geometric “ring decomposition” as opposed to layer decomposition.

that 3-SAT cannot be solved in  $O(2^{o(n)})$  time, where  $n$  denotes the number of variables of the 3-SAT formula.

## References

- [1] J. Alber, H. L. Bodlaender, H. Fernau, and R. Niedermeier. Fixed parameter algorithms for planar dominating set and related problems. In *Proc. 7th Scandinavian Workshop on Algorithm Theory SWAT*, vol. 1851 of *LNCS*, pp. 97–110. Springer, 2000.
- [2] J. Alber, H. L. Bodlaender, H. Fernau, T. Kloks and R. Niedermeier. Fixed parameter algorithms for DOMINATING SET and related problems on planar graphs. *Algorithmica*, 33:461–493, 2002.
- [3] J. Alber, F. Dorn, and R. Niedermeier. Experimental evaluation of a tree decomposition algorithm for vertex cover on planar graphs. Accepted for publication in *Discrete Applied Mathematics*, 2003.
- [4] J. Alber, H. Fan, M. Fellows, H. Fernau, R. Niedermeier, F. Rosamond, and U. Stege. Refined Search Tree Technique for Dominating Set on Planar Graphs. In *Proc. 26th International Symposium on Mathematical Foundations of Computer Science MFCS*, vol. 2136 of *LNCS*, pp. 111–122. Springer, 2001. Long version to appear in *Journal of Computer and System Sciences*.
- [5] J. Alber, M. R. Fellows, and R. Niedermeier. Efficient data reduction for DOMINATING SET: a linear problem kernel for the planar case. In *Proc. 8th Scandinavian Workshop on Algorithm Theory SWAT*, vol. 2368 of *LNCS*, pp. 150–159. Springer, 2002. Long version to appear in *Journal of the ACM*.
- [6] J. Alber, H. Fernau, and R. Niedermeier. Graph separators: a parameterized view. In *Proc. 7th Annual International Computing and Combinatorics Conference COCOON*, vol. 2108 of *LNCS*, pp. 318–327. Springer, 2001. Long version available as Technical Report WSI-2001-8, Wilhelm-Schickard-Institut für Informatik, Universität Tübingen (Germany). Most of the material appeared in *Journal of Computer and System Sciences*, 67:808–832, 2003. Some additional speed-up techniques were exhibited in “Graph separator algorithms: a refined analysis”. In *Proc. 28th Workshop on Graph-Theoretic Concepts in Computer Science WG*. vol. 2573 of *LNCS*, pp. 186–197. Springer, 2002.
- [7] J. Alber, H. Fernau and R. Niedermeier. Parameterized complexity: exponential speedup for planar graph problems. In *Proc. 15th International Colloquium on Automata, Languages and Programming ICALP*, vol. 2076 of *LNCS*, pp. 261–272. Springer, 2001.
- [8] J. Alber, J. Gramm, and R. Niedermeier. Faster exact algorithms for hard problems: A parameterized point of view. *Discrete Mathematics*, 229:3–27, 2001.



- [9] N. Alon, P. D. Seymour, and R. Thomas. A separator theorem for graphs with an excluded minor and its applications. In *Proc. 22nd ACM Symposium on the Theory of Computing STOC*, pp. 293–299, 1990.
- [10] B. S. Baker. Approximation algorithms for NP-complete problems on planar graphs. *Journal of the Association for Computing Machinery*, 41(1):153–180, 1994.
- [11] R. Bar-Yehuda and S. Even. A local-ratio theorem for approximating the weighted vertex cover problem. *Annals of Discr. Math.*, 25:27–46, 1985.
- [12] H. L. Bodlaender. Dynamic programming on graphs of bounded treewidth. In *Proc. 15th International Colloquium on Automata, Languages and Programming ICALP*, vol. 317 of *LNCS*, pp. 105–118. Springer, 1988.
- [13] H. L. Bodlaender. Treewidth: Algorithmic techniques and results. In *Proc. 22nd Mathematical Foundations of Computer Science MFCS*, vol. 1295 of *LNCS*, pp. 19–36. Springer, 1997.
- [14] H. L. Bodlaender. A partial  $k$ -arboretum of graphs with bounded treewidth. *Theoretical Computer Science*, 209:1–45, 1998.
- [15] L. Cai and D. Juedes. On the existence of subexponential parameterized algorithms. *Journal of Computer and System Sciences*, 67:789–807, 2003.
- [16] L. Cai and D. Juedes. Subexponential parameterized algorithms collapse the  $W$ -hierarchy. In *Proc. 28th International Colloquium on Automata, Languages and Programming ICALP*, vol. 2076 of *LNCS*, pp. 273–284. Springer, 2001.
- [17] J. Chen, I. Kanj, and W. Jia. Vertex cover: Further observations and further improvements. *Journal of Algorithms*, 41: 280–301, 2001.
- [18] N. Chiba, T. Nishizeki, S. Abe, and T. Ozawa. A linear algorithm for embedding planar graphs using PQ-trees. *Journal of Computer and System Sciences*, 30:54–76, 1985.
- [19] E. D. Demaine, M. T. Hajiaghayi, and D. M. Thilikos. Exponential speedup of fixed parameter algorithms on  $K_{3,3}$ -minor-free or  $K_5$ -minor-free graphs. Technical Report MIT-LCS-TR-838, 2002. Abstract version in *Proc. 13th Annual International Symposium on Algorithms and Computation ISAAC*, vol. 2518 of *LNCS*, pp. 262–273. Springer, 2002.
- [20] H. N. Djidjev and S. Venkatesan. Reduced constants for simple cycle graph separation. *Acta Informatica*, 34:231–243, 1997.
- [21] R. G. Downey. Parameterized complexity for the skeptic (invited paper). In *Proc. of 18th IEEE Conference on Computational Complexity*, July 2003.
- [22] R. G. Downey and M. R. Fellows. *Parameterized Complexity*. Springer, 1999.
- [23] R. G. Downey, M. R. Fellows, R. Niedermeier, and P. Rossmanith. Parameterized complexity. Dagstuhl-Seminar-Report No. 316, August 2001.

- [24] R. G. Downey, M. R. Fellows, and U. Stege. Parameterized complexity: A framework for systematically confronting computational intractability. *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, 49:49–99, 1999.
- [25] M. R. Fellows. Parameterized complexity: the main ideas and connections to practical computing. In *Experimental Algorithmics*, vol. 2547 of *LNCS*, pp. 51–77. Springer, 2002.
- [26] F. V. Fomin and D. M. Thilikos. A new upper bound on the decomposability of planar graphs and fixed parameter algorithms. Technical report, Departament de Llenguatges i Sistemes Informàtics, Universitat Politècnica de Catalunya, Barcelona, 2002.
- [27] F. V. Fomin and D. M. Thilikos. Dominating sets and local treewidth. In *Proc. 11th Annual European Symposium on Algorithms ESA*, vol. 2832 of *LNCS*, pp. 221–229, Springer, 2003.
- [28] F. V. Fomin and D. M. Thilikos. Dominating sets in planar graphs: branch-width and exponential speed-up. In *Proc. 14th Annual ACM-SIAM Symposium on Discrete Algorithms SODA*, pp. 168–177, 2003.
- [29] D. S. Hochbaum, editor. *Approximation algorithms for NP-hard problems*. Boston, MA: PWS Publishing Company, 1997.
- [30] A. Kanevsky. Finding all minimum-size separating vertex sets in a graph. *Networks*, 23:533–541, 1993.
- [31] I. A. Kanj and L. Perkovič. Improved parameterized algorithms for planar dominating set. In *Proc. 27th International Symposium Mathematical Foundations of Computer Science MFCS*, vol. 2420 of *LNCS*, pp. 399–410. Springer, 2002.
- [32] S. Khanna and R. Motwani. Toward a syntactic characterization of PTAS. In *Proc. 28th Annual Symposium on Theory of Computing STOC*, pp. 329–337. ACM Press, 1996.
- [33] S. Khuller. Algorithms column: the vertex cover problem. *ACM SIGACT News*, 33(2):31–33, 2002.
- [34] T. Kloks. *Treewidth: Computations and Approximations*, vol. 842 of *LNCS*. Springer, 1994.
- [35] T. Kloks, C. M. Lee, and J. Liu. New algorithms for  $k$ -face cover,  $k$ -feedback vertex set, and  $k$ -disjoint cycles on plane and planar graphs. In *Proc. 28th Workshop on Graph-Theoretic Concepts in Computer Science WG*, vol. 2537 of *LNCS*, pp. 282–295. Springer, 2002.
- [36] J. v. Leeuwen. Graph algorithms. In *Handbook of Theoretical Computer Science, Vol. A: Algorithms and Complexity*, pp. 525–631. North Holland, 1990.
- [37] R. J. Lipton and R. E. Tarjan. A separator theorem for planar graphs. *SIAM Journal of Applied Mathematics*, 36(2):177–189, 1979.

- [38] R. J. Lipton and R. E. Tarjan. Applications of a planar separator theorem. *SIAM Journal on Computing*, 9(3):615–627, 1980.
- [39] M. Mahajan and V. Raman. Parameterizing above guaranteed values: MaxSat and MaxCut. *Journal of Algorithms*, 31:335–354, 1999.
- [40] K. Mehlhorn and S. Näher. *LEDA: A Platform of Combinatorial and Geometric Computing*. Cambridge University Press, Cambridge, England, 1999.
- [41] G. L. Nemhauser and J. L. E. Trotter. Vertex packing: structural properties and algorithms. *Mathematical Programming*, 8:232–248, 1975.
- [42] R. Niedermeier and P. Rossmanith. On efficient fixed-parameter algorithms for WEIGHTED VERTEX COVER. *Journal of Algorithms*, 47(2):63–77, 2003.
- [43] V. T. Paschos. A survey of approximately optimal solutions to some covering and packing problems. *ACM Computing Surveys*, 29(2):171–209, 1997.
- [44] N. Robertson, D. P. Sanders, P. Seymour, and R. Thomas. Efficiently four-coloring planar graphs. In *Proc. 28th ACM Symposium on the Theory of Computing STOC*, pp. 571–575, 1996.
- [45] J. A. Telle. Complexity of domination-type problems in graphs. *Nordic Journal of Comp.*, 1:157–171, 1994.
- [46] J. A. Telle and A. Proskurowski. Practical algorithms on partial  $k$ -trees with an application to domination-like problems. In *Algorithms and Data Structures, Proc. 3rd WADS*, vol. 709 of *LNCS*, pp. 610–621. Springer, 1993.
- [47] M. Yannakakis and F. Gavril. Edge dominating sets in graphs. *SIAM Journal on Applied Mathematics*, 38(3):364–372, 1980.